

Generalizations of Wiener polarity index and terminal Wiener index

Aleksandar Ilić [‡]

Faculty of Sciences and Mathematics, Višegradska 33, 18 000 Niš
University of Niš, Serbia
e-mail: aleksandari@gmail.com

Milovan Ilić

Faculty of Information Technology, Trg republike 3, 11 000 Beograd
University of Belgrade, Serbia
e-mail: ilic.milovan@gmail.com

June 16, 2011

Abstract

In theoretical chemistry, distance-based molecular structure descriptors are used for modeling physical, pharmacologic, biological and other properties of chemical compounds. We introduce a generalized Wiener polarity index $W_k(G)$ as the number of unordered pairs of vertices $\{u, v\}$ of G such that the shortest distance $d(u, v)$ between u and v is k . For $k = 3$, we get standard Wiener polarity index. Furthermore, we generalize the terminal Wiener index $TW_k(G)$ as the sum of distances between all pairs of vertices of degree k . For $k = 1$, we get standard terminal Wiener index. In this paper we describe a linear time algorithm for computing these indices for trees and partial cubes, and characterize extremal trees maximizing the generalized Wiener polarity index and generalized terminal Wiener index among all trees of given order n .

Key words: Distance in graphs; Wiener polarity index; terminal Wiener index; Wiener index; Partial cube; Graph algorithm.

AMS Classifications: 05C12, 92E10.

1 Introduction

Let $G = (V, E)$ be a connected simple graph with $n = |V|$ vertices and $m = |E|$ edges. For vertices $u, v \in V$, the distance $d(u, v)$ is defined as the length of the shortest path between u and v in G . The diameter $diam(G)$ is the greatest distance between two vertices of G . Let $d_k(u)$ denotes the number of vertices on distance k from the vertex u . Let $deg(v)$ denotes the degree of the vertex v .

In theoretical chemistry molecular structure descriptors (also called topological indices) are used for modeling physico-chemical, pharmacologic, toxicologic, biological and other properties of chemical compounds [12]. There exist several types of such indices, especially those based on vertex and edge distances. Arguably the best known of these indices is the Wiener index W , defined as the sum of distances between all pairs of vertices of the molecular graph [9]

$$W(G) = \sum_{u, v \in V(G)} d(u, v).$$

Besides of use in chemistry, it was independently studied due to its relevance in social science, architecture, and graph theory. With considerable success in chemical graph theory, various extensions and generalizations of the Wiener index are recently put forward [3, 11, 24].

The Wiener polarity index of a graph G is defined as the number of unordered pairs of vertices $\{u, v\}$ of G such that the shortest distance $d(u, v)$ between u and v is 3,

$$WP(G) = |\{(u, v) \mid d(u, v) = 3, u, v \in V\}|.$$

Hosoya [14] found a physico-chemical interpretation of WP . Du, Li and Shi [8] described a linear time algorithm for computing the Wiener polarity index of trees, and characterized the trees maximizing the index among all trees of given order. Deng et al. [4, 5, 6] and Liu et al. [21] characterized extremal n -vertex trees with given diameter, number of pendent vertices or maximum vertex degree.

The terminal Wiener index of a graph G is defined by Gutman, Furtula and Petrović in [10] as the sum of distances between all pairs of pendent vertices of G

$$TW(G) = \sum_{\substack{u, v \in V(G) \\ \deg(u)=\deg(v)=1}} d(u, v).$$

Furthermore, the authors described a simple method for computing TW of trees and characterized the trees with minimum and maximum TW . Heydari and Gutman [13] provided a formula for calculating TW of thorn graphs, while Deng and Zhang [7] studied equiseparability on terminal Wiener index. Independently, Székely et al. [23] introduced the same index (the sum of distances between the leaves of a tree) and studied the correlation between various distance-based topological indices.

The paper is organized as follows. In Section 2 we introduce generalization of the Wiener polarity index and characterize the trees maximizing the generalized Wiener polarity index among all trees of given order, while in Section 3 we designed linear algorithm for calculating this index. In Section 4 we introduce generalization of the terminal Wiener index and characterize trees maximizing the generalized terminal Wiener index among all trees of given order and $k \leq 3$. In Section 5 we present formula for calculation of $TW_k(G)$ for partial cubes and in particular closed formula for TW_3 of coronene series H_k . We close the paper in Section 6 by proposing new problems for research.

2 Generalization of Wiener polarity index

For $k \geq 1$, we define the generalized Wiener polarity index as the number of unordered pairs of vertices $\{u, v\}$ of G such that the shortest distance $d(u, v)$ between u and v is k ,

$$W_k(G) = \frac{1}{2} \sum_{v \in V(G)} d_k(v).$$

Notice that $W(G) = \sum_{k=1}^{diam(G)} W_k(G)$.

For $k = 1$, it can be easily seen that $W_1(G) = m$, where m is the number of edges. For $k = 2$, we have

$$W_2(G) = \sum_{v \in V} \binom{deg(v)}{2} = \frac{\sum_{v \in V} deg^2(v)}{2} - m = M_1(G) - m,$$

where $M_1(G)$ denotes the first Zagreb index of a graph [22].

For $k = 3$ we have the Wiener polarity index,

$$\begin{aligned} W_3(T) &= \sum_{uv \in E} (deg(v) - 1)(deg(u) - 1) = \sum_{uv \in E} deg(u)deg(v) - \sum_{v \in V} deg^2 v + m \\ &= M_2(T) - M_1(T) + m, \end{aligned}$$

where $M_2(T)$ denotes the second Zagreb index of a graph [15].

In the following assume that $k \geq 3$. If the diameter of T is less than k , then $W_k(T) = 0$. Therefore, the minimum value of $W_k(T)$ is zero, and it is achieved for all trees with $diam(T) < k$ (for example the star S_n). On the other hand, we will prove that the maximum value of $W_k(T)$ is achieved for a tree with diameter k and with all pendent vertices on distance k .

The group of pendent vertices is defined as the set of all pendent vertices attached to the same unique neighbor. Let A_1 and A_2 be two different groups of pendent vertices with the unique neighbors w_1 and w_2 , such that the distance between arbitrary pendent vertices from these groups is not equal to k . Let

$$p_1 = \sum_{v \in A_1} d_k(v) \quad \text{and} \quad p_2 = \sum_{u \in A_2} d_k(u).$$

Without loss of generality assume that $p_1 \geq p_2$. If we remove all pendent vertices from A_2 and add them to the set A_1 , we get a new tree T' such that

$$W_k(T') - W_k(T) = p_1 - p_2 \geq 0.$$

By repetitive application of this transformation, we will get a new tree with possibly increased generalized Wiener polarity index. The diameter of T' is not greater than the diameter of T and each transformation introduces at most one new pendent vertex. By choosing two most distant groups of pendent vertices, we will get the extremal tree with diameter equal to k and all pendent vertices are on distance k or 2.

Assume that there are p groups of pendent vertices with sizes a_1, a_2, \dots, a_p and $a_1 + a_2 + \dots + a_p = q$. Since $diam(T) = k$, we have $n - k + 1 \geq q \geq 2$. The distance between any two pendent vertices not from the same group is equal to k , and therefore

$$W_k(T) = \frac{1}{2} \sum_{i=1}^p a_i(q - a_i) = \frac{1}{2} \left(q^2 - \sum_{i=1}^p a_i^2 \right).$$

The minimum value of $\sum_{i=1}^p a_i^2$ under the condition $\sum_{i=1}^p a_i = q$ is achieved if and only if all numbers a_i are as close as possible, i. e. $|a_i - a_j| \leq 1$ for all $1 \leq i < j \leq p$. This can be easily proved by the transformation $(a_i, a_j) \mapsto (a_i + 1, a_j - 1)$ with $a_j \geq a_i + 2$, since

$$(a_i + 1)^2 + (a_j - 1)^2 - a_i^2 - a_j^2 = 2(a_i - a_j) + 2 > 0.$$

Notice that the tree is uniquely determined by the distances between pendent vertices [26]. A starlike tree is a tree with exactly one vertex of degree at least 3. If $p = 2$, we have $W_k(T) = a_1 a_2$ and $a_1 + a_2 = n - k + 1$ and finally

$$W_k(T) = \left\lfloor \frac{n - k + 1}{2} \right\rfloor \cdot \left\lceil \frac{n - k + 1}{2} \right\rceil$$

Let $p > 2$. For k odd, we can consider two groups of pendent vertices together with the unique path connecting them. The third group of pendent vertices must be on equal distance

from both groups and that is impossible. Therefore, the extremal value for odd k is achieved for $p = 2$.

For k even, similarly it can be concluded that there is a unique tree with p groups of pendent vertices (starlike tree with p paths with equal lengths together with groups of pendent vertices attached at the pendent vertices of these p paths). For $p > 2$, we have

$$n = 1 + p \left(\frac{k}{2} - 1 \right) + \sum_{i=1}^p a_i = 1 + p \left(\frac{k}{2} - 1 \right) + q,$$

and since $p \leq q$, we have $p \leq 2 \cdot \frac{n-1}{k}$. Therefore, using Cauchy–Schwartz inequality it follows

$$\begin{aligned} W_k(T) &= \frac{1}{2} \left(q^2 - \sum_{i=1}^p a_i^2 \right) \\ &\leq \frac{1}{2} \left(q^2 - \frac{q^2}{p} \right) \\ &= \frac{1}{2} \left(n - 1 - \frac{pk}{2} + p \right)^2 \left(1 - \frac{1}{p} \right). \end{aligned}$$

Let

$$f(p) = \frac{1}{2} \left(n - 1 - \frac{pk}{2} + p \right)^2 \left(1 - \frac{1}{p} \right),$$

for $2 < p < 2 \cdot \frac{n-1}{k} < 2 \cdot \frac{n-1}{k-2}$. The first derivative equals

$$f'(p) = \frac{(2 - 2n - 2p + kp)(2 - 2n + 2p - kp - 4p^2 + 2kp^2)}{4p^2}.$$

Finally it holds that $f(p)$ is increasing for $2 < p < p^*$ and decreasing for $p^* \leq p < 2 \cdot \frac{n-1}{k}$, where

$$p^* = \frac{k - 2 + \sqrt{(k-2)(16n+k-18)}}{4(k-2)} = \frac{1}{4} + \frac{1}{4} \sqrt{\frac{16n+k-18}{k-2}}$$

is the second largest root of $f'(p) = 0$. Therefore, the maximum of the Wiener polarity index for k even is achieved for $p = 2$ or p that is near to p^* (see Figure 1).




Figure 1: The extremal tree with maximal generalized Wiener polarity index on $n = 22$ and $k = 6$.

3 Linear algorithm for W_k of trees

Let T be an arbitrary tree rooted at the vertex 1. We will process the vertices according to the distance from the root vertex or in the recursive depth first search method [2]. For each vertex v , we will keep the vector $a[v]$ of the length $k + 1$ that stores the number of vertices in the subtree under v on distances $0, 1, 2, \dots, k$. It follows that $a[v][0] = 1$ and $a[v][1] = \deg(v) - 1$ for all vertices different than the root. The matrix a with dimensions $n \times k$ is computed recursively in the first procedure.

In the second procedure we calculate the generalized Wiener polarity index. In DFS tree, for each path of the length k there is a unique vertex v on the smallest distance from the root. Therefore, we need to traverse all vertices v and count the vertices that are on distance k in the subtree under v , such that the unique path connecting these vertices contains v . In the first case, we just add $2a[v][k]$. In the second case, we need to consider all neighbors u of v different than $\text{parent}[v]$ and for each $j = 0, 1, \dots, k - 2$ count the number of vertex pairs (x, y) such that:

- x is in the subtree under u and y is not;
- x is on the distance j from u ;
- y is on the distance $k - j - 1$ from v and under v .

Finally we counted every vertex pair on distance k twice, as showed in the second procedure.

Procedure DFS(vertex v)

Input: The adjacency list of the tree T with the root vertex $root$.

Output: The array parent and matrix a .

$a[v][0] = 1;$

foreach neighbor u of v **do**

if ($\text{parent}[u] = 0$) and ($u \neq \text{root}$) **then**

$\text{parent}[u] = v;$

 DFS(u);

for $i = 0$ **to** $k - 1$ **do**

$a[v][i + 1] = a[v][i + 1] + a[u][i];$

end

end

end

4 Generalization of terminal Wiener index

For $k \geq 1$, we define the generalized terminal Wiener index as the sum of the distances between all pairs of its vertices of degrees k ,

$$TW_k(G) = \sum_{\substack{u, v \in V(G) \\ \deg(u) = \deg(v) = k}} d(u, v).$$

For k regular graphs, $WT_k(G) = W(G)$ and $WT_i(G) = 0$ for $i \neq k$. For $k = 1$, we have terminal Wiener index.

Procedure GWP

Input: The adjacency list of the tree T with the root vertex $root$ and the array a .

Output: The generalized Wiener polarity index Wk .

$Wk = 0$;

for $v = 1$ **to** n **do**

$Wk = Wk + 2 \cdot a[v][k]$;

foreach neighbor u of v **do**

if $parent[v] \neq u$ **then**

for $i = 0$ **to** $k - 2$ **do**

$Wk = Wk + a[u][i] \cdot (a[v][k - 1 - i] - a[u][k - 2 - i])$;

end

end

end

end

return $Wk/2$;

Theorem 4.1 ([9]) *Let T be a tree on n vertices. Then,*

$$(n - 1)^2 = W(S_n) \leq W(T) \leq W(P_n) = \binom{n + 1}{3},$$

with equality if and only if $T \cong S_n$ or $T \cong P_n$.

Theorem 4.2 ([16]) *Let w be a vertex of a nontrivial connected graph G . For nonnegative integers p and q , let $G(p, q)$ denote the graph obtained from G by attaching to vertex w pendent paths $P = wv_1v_2 \dots v_p$ and $Q = wu_1u_2 \dots u_q$ of lengths p and q , respectively. If $p \geq q \geq 1$, then*

$$W(G(p, q)) < W(G(p + 1, q - 1)).$$

For $k = 1$, Gutman, Furtula and Petrović in [10] characterized the extremal trees within the class of all trees with n vertices that maximize terminal Wiener index. For $k = 2$, the unique tree that maximizes TW_2 is the path P_n . Interestingly, the path P_n is the unique tree that minimizes TW_1 . This follows from Theorems 4.1 and 4.2 and the simple fact that in every tree there are at least 2 pendent vertices.

Let T^* be the extremal tree that maximizes the generalized terminal Wiener index for $k \geq 3$. It can be easily proved that there are no pendent paths $P = v_1v_2 \dots v_p$ of length greater than 2 attached at some vertex of T^* . Otherwise, remove pendent edges one by one and subdivide any edge e such that both components of $T^* - e$ contain vertices of degree k – this way we increase TW_k . If such edge does not exist, remove the edge $v_{p-1}v_p$ and add new edge $v_{p-2}v_p$ – this way the generalized terminal Wiener index remains the same, or increases if and only if $k = 3$.

Lemma 4.3 *Among trees on n vertices the maximal number of vertices of degree $k \geq 2$ is*

$$m(n, k) = \left\lfloor \frac{n - 2}{k - 1} \right\rfloor.$$

Proof. Consider the induced graph H composed of the vertices of degree k . Let h be the number of vertices in H , and let f be the number of edges in H . Since H is acyclic and

possible disconnected, we have $f \leq h - 1$. Furthermore, for the number of edges in tree T holds

$$h \cdot k - (h - 1) \leq h \cdot k - f \leq n - 1.$$

It follows that $h(k - 1) \leq n - 2$, which completes the proof. \blacksquare




Figure 2: The extremal tree with maximal generalized terminal Wiener index on $n = 20$ and $k = 4$.

The $C_{n,k,p}$ be the caterpillar obtained from a path of length $s + 2 = n - p \cdot (k - 2)$, by attaching $k - 2$ pendent vertices to exactly p vertices of a path $P_{s+2} = v_0 v_1 \dots v_s v_{s+1}$, starting from the vertices v_1 and v_s on the both ends towards center symmetrically (see Figure 2). The generalized terminal Wiener index can be easily calculated,

$$\begin{aligned} TW_k(C_{n,k,p}) &= \sum_{\substack{1 \leq i \leq \lfloor p/2 \rfloor \\ s - \lfloor p/2 \rfloor + 1 \leq j \leq s}} d(v_i, v_j) \\ &= (p - 1)(s - 1) + (p - 3)(s - 3) + (p - 5)(s - 5) + \dots \\ &= \begin{cases} \frac{1}{12}p(3ps - p^2 - 2), & \text{if } p \text{ is even} \\ \frac{1}{12}(p + 1)(p - 1)(3s - p), & \text{if } p \text{ is odd.} \end{cases} \\ &= \begin{cases} \frac{1}{12}p(3np + 5p^2 - 3kp^2 - 2 - 6p), & \text{if } p \text{ is even} \\ \frac{1}{12}(p + 1)(p - 1)(3n + 5p - 3kp - 6), & \text{if } p \text{ is odd.} \end{cases} \end{aligned} \quad (1)$$




Figure 3: Transformation that increases generalized terminal Wiener index TW_3 .

We call a caterpillar C 3-bounded if all vertices of C have degree less than or equal to 3.

Theorem 4.4 Let T be a tree on $n > 4$ vertices. Then

$$TW_3(T) \leq TW_3(C_{n,3,\lfloor n/2 \rfloor - 1}),$$

with equality if and only if $T \cong C_{n,3,\lfloor n/2 \rfloor - 1}$.

Proof. Let T^* be a rooted tree with maximal value of generalized terminal Wiener index for $k = 3$. If T^* is not a caterpillar, consider a branching vertex w such that in the subtree under

w there are only 3-bounded caterpillars attached at w . Let C_1 and C_2 be two caterpillars attached at w , such that C_1 has p vertices v_1, v_2, \dots, v_p of degree 3 and C_2 has q vertices u_1, u_2, \dots, u_q of degree 3 (see Figure 3). Without loss of generality, we can assume that the number of vertices of degree 3 in $G = T^* \setminus \{C_1, C_2\}$ is greater than or equal to p and q , namely $r \geq p \geq q$.

Let a_1, a_2, \dots, a_p be the distances from the vertex w to the vertices v_1, v_2, \dots, v_p and b_1, b_2, \dots, b_q be the distance from the vertex w to the vertices u_1, u_2, \dots, u_q . Let $D(w)$ be the sum of distances from the vertex w to the vertices of degree 3 in the subgraph G .

The generalized terminal Wiener index of the tree T^* equals

$$\begin{aligned} TW_3(T^*) &= TW_3(G) + (p+q)D(w) + r \left(\sum_{i=1}^p a_i + \sum_{j=1}^q b_j \right) \\ &\quad + \sum_{i < j} (a_j - a_i) + \sum_{i < j} (b_j - b_i) + q \cdot \sum_{i=1}^p a_i + p \cdot \sum_{j=1}^q b_j \end{aligned}$$

After reattaching the caterpillar C_2 to the end of caterpillar C_1 , the degree of vertex w remains the same. Furthermore, the vertex v_p has degree 3 and the generalized terminal Wiener index of transformed tree T' equals

$$\begin{aligned} TW_3(T') &= TW_3(G) + (p+q)D(w) + r \cdot \sum_{i=1}^p a_i + r \left(q \cdot a_p + \sum_{j=1}^q b_j \right) \\ &\quad + \sum_{i < j} (a_j - a_i) + \sum_{i < j} (b_j - b_i) + q \cdot \left(p \cdot a_p - \sum_{i=1}^p a_i \right) + p \cdot \sum_{j=1}^q b_j \end{aligned}$$

By subtraction, we get

$$TW_3(T') - TW_3(T^*) = rq \cdot a_p + pq \cdot a_p - 2q \cdot \sum_{i=1}^p a_i,$$

Since $r \geq p$ and $a_p \geq a_i$ for $i = 1, 2, \dots, p$, we have

$$\begin{aligned} TW_3(T') - TW_3(T^*) &= q \left(a_p(r+p) - 2 \sum_{i=1}^p a_i \right) \\ &\geq 2q \cdot \sum_{i=1}^p (a_p - a_i) \geq 0, \end{aligned}$$

with equality if and only if $p = q = 1$.

It follows that using this transformation we can not decrease the generalized Wiener index TW_3 . If $\deg(w) > 3$, we can remove the pendent edge and subdivide some edge e such that both components of $T^* - e$ contain vertices of degree 3, and further increase the generalized Wiener index. Finally, we conclude that the extremal tree is a caterpillar of the form $C_{n,k,p}$.

For $k = 3$, from (1) we have

$$f(p) = \begin{cases} \frac{1}{12}p(3np - 4p^2 - 2 - 6p), & \text{if } p \text{ is even} \\ \frac{1}{12}(p+1)(p-1)(3n - 4p - 6), & \text{if } p \text{ is odd.} \end{cases}$$

and

$$f(p) - f(p-2) = \begin{cases} (p-1)(n-2p) - 1, & \text{if } p \text{ is even} \\ (p-1)(n-2p), & \text{if } p \text{ is odd,} \end{cases}$$

which is greater than zero since $n \geq 2p + 2$. By direct verification, we get $f(\lfloor \frac{n}{2} \rfloor - 1) > f(\lfloor \frac{n}{2} \rfloor - 2)$ for $n > 4$.

Therefore, it holds

$$TW_3(T) \leq TW_3(C_{n,3,\lfloor n/2 \rfloor - 1}),$$

with equality if and only if $T \cong C_{n,3,\lfloor n/2 \rfloor - 1}$. ■

5 Calculating TW_k of partial cubes

Let G be a connected graph. Then $e = xy$ and $f = uv$ are in the Djoković-Winkler Θ relation [25] if

$$d(x, u) + d(y, v) \neq d(x, v) + d(y, u).$$

The relation Θ is always reflexive and symmetric, and is transitive on partial cubes. Therefore, Θ partitions the edge set of a partial cube G into equivalence classes F_1, F_2, \dots, F_s , called Θ -classes. For any $1 \leq i \leq s$, the graph $G - F_i$ consists of two connected components. The vertex sets of these components will be denoted with $W_{i,0}$ and $W_{i,1}$, because they can be described as the vertices whose i -th coordinate is 0 and 1, respectively. The sets $W_{i,\chi}$, $1 \leq i \leq s$, $\chi \in \{0, 1\}$ are called halfspaces of G , while $W_{i,0}$ and $W_{i,1}$ are complementary halfspaces [18] and it holds $|W_{(i,0)}|^{(k)} + |W_{(i,1)}|^{(k)} = 1$.

Let G be a partial cube with halfspaces $W_{i,\chi}$, $1 \leq i \leq s$, $\chi \in \{0, 1\}$. For any $1 \leq i \leq s$ and any $\chi \in \{0, 1\}$, $|W_{(i,\chi)}|^{(k)}$ denotes the number of vertices with degree k .

Theorem 5.1 *Let G be a partial cube with halfspaces $W_{i,\chi}$, $1 \leq i \leq s$, $\chi \in \{0, 1\}$. Then*

$$TW_k(G) = \sum_{i=1}^s |W_{(i,0)}|^{(k)} \cdot |W_{(i,1)}|^{(k)}.$$

Proof. G is a partial cube, hence vertices of G can be considered as a binary s -tuple $u = u_1 u_2 \dots u_s$. Moreover, since G is isometric in Q_s , the distance between two vertices is the number of positions in which they differ. Set $\delta(x, y) = 0$ if $x = y$, and $\delta(x, y) = 1$ for $x \neq y$. Then

$$\begin{aligned} TW_k(G) &= \sum_{\substack{u, v \in V(G) \\ \deg(u) = \deg(v) = k}} d(u, v) \\ &= \sum_{\substack{u, v \in V(G) \\ \deg(u) = \deg(v) = k}} \sum_{i=1}^s \delta(u_i, v_i) \\ &= \sum_{i=1}^s \left(\sum_{\substack{u, v \in V(G) \\ \deg(u) = \deg(v) = k}} \delta(u_i, v_i) \right) \\ &= \sum_{i=1}^s |W_{(i,0)}|^{(k)} \cdot |W_{(i,1)}|^{(k)}. \end{aligned}$$

This completes the proof. ■

An advantage of Theorem 5.1 comparing to computing $TW_k(G)$ by the definition is that we do not need to compute distances, but only to count vertices in the classes. This theorem

can be considered as another instance of Klavžar cut method. For its general description and an overview of its applications in chemical graph theory see survey [19].

As an example, we obtain a closed expression for TW_3 of the coronene/circumcoronene homologous series H_k . In Figure 4, $2k - 1$ horizontal elementary cuts of H_k are presented. There exist two additional groups of $2k - 1$ equivalent cuts, obtained by rotating the former group by $\frac{\pi}{3}$ and $-\frac{\pi}{3}$. The number of vertices of H_k equals $n_k = 6k^2$, while there are exactly $6k$ vertices of degree two.




Figure 4: The Coronene / Circumcoronene H_3 .

Using symmetry, the contribution of the elementary cut C_i is equal to the contribution of C_{2k-i} , $i = 1, 2, \dots, k - 1$. By induction it follows that for $i = 1, 2, \dots, k$ the number of vertices above cut C_i equals $i(2k + i)$, while the number of vertices of degree 2 equals $k + 2i$. Therefore,

$$\begin{aligned} \frac{1}{3}TW_3(H_k) &= (3k^2 - 3k)^2 + 2 \sum_{i=1}^{k-1} (2ki + i^2 - k - 2i)(6k^2 - 6k - 2ki - i^2 + k + 2i) \\ &= \frac{164k^5}{15} - \frac{82k^4}{3} + \frac{58k^3}{3} - \frac{5k^2}{3} - \frac{19k}{15}. \end{aligned}$$

Finally, we derive the fifth-order polynomial formula for the generalized terminal Wiener index of H_k

$$TW_3(H_k) = \frac{1}{5}(k - 1)k(2k - 1)(82k^2 - 82k - 19).$$

Using similar methods as in [1, 17, 20] one can obtain the closed formulas for other chemical graphs (trees, benzenoid chains, phenylenes, ...) and design a linear algorithm for TW_k of benzenoid systems.

6 Concluding remarks

The Wiener polarity index and the terminal Wiener index are very new molecular-structure descriptors and only a limited number of mathematical and chemical properties were established so far. In this paper we generalized these indices and open new perspectives for the future research.

Another generalization of these indices may be the following

$$W_k^*(G) = |\{(u, v) \mid d(u, v) \leq k, u, v \in V\}| = W_1(G) + W_2(G) + \dots + W_k(G)$$

and

$$TW_k^*(G) = \sum_{\substack{u,v \in V(G) \\ \deg(u) \leq k, \deg(v) \leq k}} d(u,v) \geq TW_1(G) + TW_2(G) + \dots + TW_k(G).$$

It would be nice to study mathematical and algorithmic properties of these indices and report their chemical relevance. These indices are obtained from the famous Wiener index, which has many applications in chemistry, graph theory and computer science.

Acknowledgement. This work was supported by Research Grant 144007 of Serbian Ministry of Science and Technological Development.

References

- [1] V. Chepoi, S. Klavžar, *The Wiener index and the Szeged index of benzenoid systems in linear time*, J. Chem. Inf. Comput. Sci. 37 (1997) 752–755.
- [2] T. H. Cormen, C. E. Leiserson, R. L. Rivest, C. Stein, *Introduction to Algorithms*, Second Edition, MIT Press, 2001.
- [3] P. Dankelmann, I. Gutman, S. Mukwembi, H. C. Swart, *The edge-Wiener index of a graph*, Discrete Math. 309 (2009) 3452–3457.
- [4] H. Deng, *On the extremal Wiener polarity index of chemical trees*, MATCH Commun. Math. Comput. Chem., in press.
- [5] H. Deng, H. Xiao, F. Tang, *The maximum Wiener polarity index of trees with k pendants*, Appl. Math. Letters 23 (2010) 710–715.
- [6] H. Deng, H. Xiao, F. Tang, *On the extremal Wiener polarity index of trees with a given diameter*, MATCH Commun. Math. Comput. Chem. 63 (2010) 257–264.
- [7] X. Deng, J. Zhang, *Equiseparability on terminal Wiener index*, in: A. V. Goldberg, Y. Zhou (Eds.), *Algorithmic Aspects in Information and Management*, Springer-Verlag, Berlin, 2009, 166–174.
- [8] W. Du, X. Li, Y. Shi, *Algorithms and extremal problem on Wiener polarity index*, MATCH Commun. Math. Comput. Chem. 62 (2009) 235–244.
- [9] A. A. Dobrynin, R. C. Entringer, I. Gutman, *Wiener index of trees: theory and applications*, Acta Appl. Math. 66 (2001) 211–249.
- [10] I. Gutman, B. Furtula, M. Petrović, *Terminal Wiener index*, J. Math. Chem. 46 (2009) 522–531.
- [11] I. Gutman, D. Vukičević, J. Žerovnik, *A class of modified Wiener indices*, Croat. Chem. Acta 77 (2004) 103–109.
- [12] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer – Verlag, Berlin, 1988.
- [13] A. Heydari, I. Gutman, *On the terminal Wiener index of thorn graphs*, Kragujevac J. Sci. 32 (2010) 57–64.

- [14] H. Hosoya, *Mathematical and chemical analysis of Wiener's polarity number*, in: D. H. Rouvray, R. B. King (Eds.), *Topology in Chemistry – Discrete Mathematics of Molecules*, Horwood, Chichester, 2002.
- [15] A. Ilić, D. Stevanović, *On Comparing Zagreb Indices*, MATCH Commun. Math. Comput. Chem. 62 (2009) 681–687.
- [16] A. Ilić, A. Ilić, D. Stevanović, *On the Wiener index and Laplacian coefficients of graphs with given diameter or radius*, MATCH Commun. Math. Comput. Chem. 63 (2010) 91–100.
- [17] A. Ilić, S. Klavžar, D. Stevanović, *Calculating the Degree Distance of Partial Hamming Graphs*, MATCH Commun. Math. Comput. Chem. 63 (2010) 411–424.
- [18] W. Imrich, S. Klavžar, *Product Graphs: Structure and Recognition*, John Wiley & Sons, New York, 2000.
- [19] S. Klavžar, *Bird's eye view of the cut method and a survey of its applications in chemical graph theory*, MATCH Commun. Math. Comput. Chem. 60 (2008) 255–274.
- [20] S. Klavžar, I. Gutman, *Wiener number of vertex-weighted graphs and a chemical application*, Discrete Appl. Math. 80 (1997) 73–81.
- [21] B. Liu, H. Hou, Y. Huang, *On the Wiener polarity index of trees with maximum degree or given number of leaves*, Comp. Math. Appl. 60 (2010) 2053–2057.
- [22] S. Nikolić, G. Kovačević, A. Milićević, N. Trinajstić, *The Zagreb indices 30 years after*, Croat. Chem. Acta 76 (2003) 113–124.
- [23] L. A. Székely, H. Wang, T. Wu, *The sum of the distances between the leaves of a tree and the 'semi-regular' property*, Discrete Math. (2010), doi:10.1016/j.disc.2010.06.005.
- [24] R. Todeschini, V. Consonni, *Handbook of Molecular Descriptors*, Wiley – VCH, Weinheim, 2000.
- [25] P. Winkler, *Isometric embeddings in products of complete graphs*, Discrete Appl. Math. 7 (1984) 221–225.
- [26] K. A. Zaretskii, *Construction of trees using the distances between pendent vertices*, Uspekhi Math. Nauk. 20 (1965) 90–92.